

Vaccination Involving the Sinovac-CoronaVac or Sinopharm COVID-19 Vaccines for Mixed Vaccine Combinations and Recovered Persons

The table lists the vaccination combinations in order to be considered fully vaccinated, for persons who:

- (a) Have received a mixed vaccine combination incorporating Sinovac-CoronaVac or Sinopharm vaccines, or,
- (b) Have recovered from COVID-19 and took the Sinovac-CoronaVac or Sinopharm vaccines.

The table also provides recommendations on the intervals between vaccine doses/recovery.

Note:

- **S** refers to either the Sinovac-CoronaVac or Sinopharm vaccine
- **W** refers to a World Health Organization's Emergency Use Listing (WHO-EUL) vaccine excluding Sinovac-CoronaVac and Sinopharm vaccines, and the Johnson & Johnson/Janssen single-dose COVID-19 vaccine.

S/N	Three-Dose Mixed Combinations Considered to be Fully Vaccinated	
1.	S-W-W W-S-W	The interval between the two W doses should be that recommended for the W vaccine or greater.
2.	S-S-W S-S-S	<p>The minimum interval between the first and second doses should be:</p> <ul style="list-style-type: none"> • 13 days for Sinovac-CoronaVac vaccine, and • 17 days for the Sinopharm vaccine. <p>The interval between the second and third doses should be at least 86 days.</p>
3.	W-S-S	<p>The minimum interval between the first and second doses is to be no less than 17 days.</p> <p>The minimum interval between the second and third doses should be:</p> <ul style="list-style-type: none"> • 17 days if the second dose is Sinovac-Coronavac vaccine, and • 17 days, if the second dose is Sinopharm vaccine.
4.	S-W-S	The minimum interval between the first and second doses is to be no less than 17 days.

		The interval between the second and third doses is to be no less than 17 days.
	Recovered Persons Vaccinated with Sinovac-CoronaVac or Sinopharm Vaccines	
4.	S-Recovered-W S-S-Recovered-W	The minimum interval between the date of COVID-19 diagnosis and the dose of W vaccine thereafter should be no less than 28 days.
5.	Recovered-S-S S-Recovered-S-S W-Recovered-S-S	The minimum interval between the date of COVID-19 diagnosis and the dose of first S vaccine thereafter should be no less than 28 days. The interval between the two S-S vaccines after having recovered should be: <ul style="list-style-type: none"> • 13 days for Sinovac-CoronaVac vaccine, and • 17 days for the Sinopharm vaccine.
6.	S-S-Recovered-S	The minimum interval between the date of COVID-19 diagnosis and the dose of the S vaccine thereafter should be no less than 28 days.
7.	S-W-Recovered W-S-Recovered	The minimum interval between the first and second doses is to be no less than 17 days. No further doses are required after having recovered for this combination of S-W vaccines ¹ .

Note: The table presents the minimum intervals accepted. The National Vaccination Programme may adopt the recommended intervals instead for better outcomes. These intervals may be longer.

¹ However, this does not apply to recovered persons who have received one dose of S or W vaccine or 2 doses of S vaccines. For this group, they will require further dose(s) of vaccine.